



# Discussion Guide

When you are considering or starting **TECENTRIQ** for metastatic non-small cell lung cancer, or mNSCLC, you or your caregiver may have questions. **Use this guide to help you prepare for speaking with your oncologist or other members of your healthcare team. This guide is not intended to replace a conversation with your doctor.**

**Start by getting to know your healthcare team.** Use the space below to fill in the names and contact information of some of the healthcare providers that may be involved in your care.

## MY HEALTHCARE TEAM

**Oncologist:** \_\_\_\_\_ **Social Worker:** \_\_\_\_\_ **Nutritionist:** \_\_\_\_\_

Phone/Email: \_\_\_\_\_ Phone/Email: \_\_\_\_\_ Phone/Email: \_\_\_\_\_

**Nurse:** \_\_\_\_\_ **Pharmacist:** \_\_\_\_\_ **Other:** \_\_\_\_\_

Phone/Email: \_\_\_\_\_ Phone/Email: \_\_\_\_\_ Phone/Email: \_\_\_\_\_

**Other Doctor:** \_\_\_\_\_ **Patient Navigator:** \_\_\_\_\_

Phone/Email: \_\_\_\_\_ Phone/Email: \_\_\_\_\_



## WHAT IS TECENTRIQ?

TECENTRIQ is a prescription medicine used to treat adults with a **type of lung cancer called non-small cell lung cancer (NSCLC)**. **TECENTRIQ may be used alone as your first treatment when your lung cancer** has spread or grown, **and** your cancer tests positive for “high PD-L1”, **and** your tumor does not have an abnormal “EGFR” or “ALK” gene.

**TECENTRIQ may be used with the medicines bevacizumab, paclitaxel, and carboplatin as your first treatment when your lung cancer** has spread or grown, **and** is a type called “non-squamous NSCLC”, **and** your tumor does not have an abnormal “EGFR” or “ALK” gene. **TECENTRIQ may be used with the medicines paclitaxel protein-bound and carboplatin as your first treatment when your lung cancer** has spread or grown, **and** is a type called “non-squamous NSCLC”, **and** your tumor does not have an abnormal “EGFR” or “ALK” gene.

**TECENTRIQ may be used alone when your lung cancer** has spread or grown, **and** you have tried chemotherapy that contains platinum, and it did not work or is no longer working. If your tumor has an abnormal “EGFR” or “ALK” gene, you should have also tried an FDA-approved therapy for tumors with these abnormal genes, and it did not work or is no longer working.

It is not known if TECENTRIQ is safe and effective when used in children for the treatment of NSCLC.

## SELECT IMPORTANT SAFETY INFORMATION

TECENTRIQ can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during your treatment or even after your treatment has ended.

Please see page 4 for more information.

**Consider asking your oncologist or their team the following questions:**

Use the space for notes on the next page to write down additional information.

**TREATING mNSCLC WITH TECENTRIQ**

- What should I know about my mNSCLC?
- Is TECENTRIQ a treatment option for my type of lung cancer?
- How does TECENTRIQ work?

TECENTRIQ may be given as a first treatment option or after other treatments have stopped working.

**QUESTIONS IF TECENTRIQ IS MY FIRST TREATMENT**

- What are the other medicines I will receive with TECENTRIQ? How often will I receive them?
- How long will I be on the other medicines?
- What side effects might I expect while on TECENTRIQ with other medicines? What do I do if I experience any?

**QUESTIONS WHEN TAKING TECENTRIQ AFTER PRIOR TREATMENTS**

- How long will I be on TECENTRIQ and how often will I receive it?
- What side effects might I expect while on TECENTRIQ? What do I do if I experience any?

**PREPARING FOR MY ONCOLOGY APPOINTMENTS AND TECENTRIQ INFUSIONS**

- How much time should I allot for my first oncology appointment? Following appointments?
- Will my oncology appointments and infusions be on the same days?
- What is my infusion schedule for TECENTRIQ?
- Can you walk me through what happens during my first infusion and following infusions?
- Should I bring someone with me when I am getting my infusion?
- What are some helpful items to have during my infusion?

**GETTING THE RIGHT SUPPORT**

- How will this treatment affect my day-to-day activities?
- I have concerns about the following, where can I go to get more information? *(check all that apply)*
  - Nutritional support
  - Emotional support
  - Getting to treatment facilities
  - Paying for treatment
  - Other

Genentech is committed to helping you get the TECENTRIQ your doctor prescribed. To learn more about Genentech Access Solutions, call **1 (877) GENENTECH (1-877-436-3683)**, or visit [Genentech-Access.com](https://www.genentech.com/access).



## WHAT IS TECENTRIQ?

TECENTRIQ is a prescription medicine used to treat adults with a **type of lung cancer called non-small cell lung cancer (NSCLC)**. **TECENTRIQ may be used alone as your first treatment when your lung cancer** has spread or grown, **and** your cancer tests positive for “high PD-L1”, **and** your tumor does not have an abnormal “EGFR” or “ALK” gene.

**TECENTRIQ may be used with the medicines bevacizumab, paclitaxel, and carboplatin as your first treatment when your lung cancer** has spread or grown, **and** is a type called “non-squamous NSCLC”, **and** your tumor does not have an abnormal “EGFR” or “ALK” gene. **TECENTRIQ may be used with the medicines paclitaxel protein-bound and carboplatin as your first treatment when your lung cancer** has spread or grown, **and** is a type called “non-squamous NSCLC”, **and** your tumor does not have an abnormal “EGFR” or “ALK” gene.

**TECENTRIQ may be used alone when your lung cancer** has spread or grown, **and** you have tried chemotherapy that contains platinum, and it did not work or is no longer working. If your tumor has an abnormal “EGFR” or “ALK” gene, you should have also tried an FDA-approved therapy for tumors with these abnormal genes, and it did not work or is no longer working.

It is not known if TECENTRIQ is safe and effective when used in children for the treatment of NSCLC.

## IMPORTANT SAFETY INFORMATION

### What is the most important information about TECENTRIQ?

TECENTRIQ can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during your treatment or even after your treatment has ended.

**Call or see your healthcare provider right away if you develop any new or worse signs or symptoms including:**

#### Lung problems

- cough
- shortness of breath
- chest pain

#### Intestinal problems

- diarrhea (loose stools) or more frequent bowel movements than usual
- stools that are black, tarry, sticky, or have blood or mucus
- severe stomach-area (abdomen) pain or tenderness

#### Liver problems

- yellowing of your skin or the whites of your eyes
- severe nausea or vomiting
- pain on the right side of your stomach area (abdomen)
- dark urine (tea colored)
- bleeding or bruising more easily than normal

#### Hormone gland problems

- headaches that will not go away or unusual headaches
- eye sensitivity to light
- eye problems
- rapid heartbeat
- increased sweating
- extreme tiredness
- weight gain or weight loss
- feeling more hungry or thirsty than usual
- urinating more often than usual
- hair loss
- feeling cold
- constipation
- your voice gets deeper
- dizziness or fainting
- changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness

Continued on page 5.

## IMPORTANT SAFETY INFORMATION CONT.

### Kidney problems

- decrease in your amount of urine
- blood in your urine
- swelling of your ankles
- loss of appetite

### Skin problems

- rash
- itching
- skin blistering or peeling
- painful sores or ulcers in mouth or nose, throat, or genital area
- fever or flu-like symptoms
- swollen lymph nodes

### Problems can also happen in other organs.

**These are not all of the signs and symptoms of immune system problems that can happen with TECENTRIQ. Call or see your healthcare provider right away for any new or worse signs or symptoms, including:**

- Chest pain, irregular heartbeat, shortness of breath, or swelling of ankles
- Confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight
- Persistent or severe muscle pain or weakness, muscle cramps
- Low red blood cells, bruising

**Infusion reactions that can sometimes be severe or life-threatening. Signs and symptoms of infusion reactions may include:**

- chills or shaking
- itching or rash
- flushing
- shortness of breath or wheezing
- dizziness
- feeling like passing out
- fever
- back or neck pain

**Complications, including graft-versus-host disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic).** These complications can be serious and can lead to death.

These complications may happen if you underwent transplantation either before or after being treated with TECENTRIQ. Your healthcare provider will monitor you for these complications.

**Getting medical treatment right away may help keep these problems from becoming more serious.** Your healthcare provider will check you for these problems during your treatment with TECENTRIQ. Your healthcare provider may treat you with corticosteroid or hormone replacement medicines. Your healthcare provider may also need to delay or completely stop treatment with TECENTRIQ if you have severe side effects.

Continued on page 6.

## IMPORTANT SAFETY INFORMATION CONT.

**Before you receive TECENTRIQ, tell your healthcare provider about all of your medical conditions, including if you:**

- have immune system problems such as Crohn's disease, ulcerative colitis, or lupus
- have received an organ transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have received radiation treatment to your chest area
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. TECENTRIQ can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with TECENTRIQ. **Females who are able to become pregnant:**
  - Your healthcare provider should do a pregnancy test before you start treatment with TECENTRIQ.
  - You should use an effective method of birth control during your treatment and for at least 5 months after the last dose of TECENTRIQ.
- are breastfeeding or plan to breastfeed. It is not known if TECENTRIQ passes into your breast milk. Do not breastfeed during treatment and for at least 5 months after the last dose of TECENTRIQ.

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**The most common side effects of TECENTRIQ when used alone include:**

- feeling tired or weak
- decreased appetite
- nausea
- cough
- shortness of breath

**The most common side effects of TECENTRIQ when used in lung cancer with other anti-cancer medicines include:**

- feeling tired or weak
- nausea
- hair loss
- constipation
- diarrhea
- decreased appetite

TECENTRIQ may cause fertility problems in females, which may affect the ability to have children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of TECENTRIQ. Ask your healthcare provider or pharmacist for more information about the benefits and side effects of TECENTRIQ.

You may report side effects to the FDA at 1-800-FDA-1088 or [www.fda.gov/safety/medwatch-fda-safety-information-and-adverse-event-reporting-program](http://www.fda.gov/safety/medwatch-fda-safety-information-and-adverse-event-reporting-program). You may also report side effects to Genentech at 1-888-835-2555.

**Please see full Prescribing Information and Medication Guide for additional Important Safety Information.**